

Crystal chemical and geochemical features of genetically different paleozoic dolomites in the Volga-Ural region

Kuznetsov V., Morozov V., Eskin A., Kolchugin A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Pleiades Publishing, Inc. The comparative analysis of primary sedimentary and secondary catagenetic dolomites revealed that they differ in many typomorphic properties. Sedimentary dolomites are characterized by the pelitomorphic and fine-grained structure, high stoichiometry, and presence of the electron-hole (p-n) centers [SO₂]⁻ and [SO₃]⁻ in the structure. Secondary dolomites have a fine- to medium-grained structure and distinguished by a lower degree of stoichiometry. Their crystal lattice shows isomorphous replacements Mg ↔ Fe and Mg ↔ Ca. They can be divided into two (massive and porous) varieties. The massive varieties are characterized by the conformal structure and xenomorphic habitus of the closely packed idiomorphic and hypidiomorphic grains. The Ca-Mg and Fe-Mg isomorphism in the massive dolomites is less expressed than in the porous varieties. These specific features of dolomite varieties are related to their different formation settings. The porous dolomites were formed in a free environment that fostered slow growth of the relatively large equant crystals with a high degree of isomorphism, whereas the massive dolomites were formed under “stringent constraints” during a faster growth of crystals.

<http://dx.doi.org/10.1134/S0024490217030038>

References

- [1] Belonin, M.D., Belonovskaya, L.G., Bulach, M.Kh., et al., Karbonatnye porody-kollektory fanerozoia neftegazonosnykh basseinov Rossii i sopredel'nykh territorii (Phanerozoic Carbonate Rock-Reservoirs in Petroliferous Basins of Russia and Adjacent Regions), St. Petersburg: Nedra, 2005, part 1.
- [2] Brannnd, U. and Veizer, J., Chemical diagenesis of a multicomponent carbonate system—1. Trace elements, J. Sedim. Petrol., 1980, vol. 50, no. 4, pp. 1219–1236.
- [3] Demchuk, I.G., Quantitative determination of isomorphic Fe in ankerites and Fe-dolomites by the X-ray structural method, Dokl. Akad. Nauk SSSR, 1985, vol. 283, no. 4, pp. 981–984.
- [4] Drits, V.A., Sakharov, B., and Milliken, K.L., New insight into structural and compositional variability in some ancient excess-Ca dolomite, Can. Mineral., 2005, vol. 43, pp. 1255–1290.
- [5] Glikin, A.E., Polimineral'no-metasomaticheskii kristallogenez (Polymineral-Metasomatic Crystallogenesis), St. Petersburg: Neva, 2004.
- [6] Gorokhov, I.M., Semikhatov, M.A., Baskakov, A.V., et al., The Sr isotopic composition in the Riphean, Vendian, and Lower Cambrian carbonate rocks, Stratigr. Geol. Korrel., 1995, vol. 3, pp. 3–33.
- [7] Kolchugin, A.N., Morozov, V.P., and Korolev, E.A., Diagenesis of carboniferous carbonate rocks reservoirs: Case study (central part of Volga-Ural basin), World Appl. Sci. J., 2013, no. 24 (6), pp. 712–718.
- [8] Kunts, A.F., Gidrotermal'no-metasomaticheskoe rudoobrazovanie v karbonatnykh porodakh (eksperimental'nye modeli i ikh prilozheniya) (Hydrothermal-Metasomatic Ore Formation in Carbonate Rocks: Experimental Models and Their Implication), Yekaterinburg: UrO RAN, 2002.

- [9] Lambert, L., Durllet, C., Loreau, J-P., and Marnier, G, Burial dissolution of micrite in Middle East carbonate reservoirs (Jurassic–Cretaceous): keys for recognition and timing, *Mar. Petr. Geol.*, 2006, no. 23, pp. 79–92.
- [10] Morozov, V.P., Korolev, E.A., and Kolchugin, A.N, Mineral-lithological criteria for the identification of zones of recent and ancient water–oil contacts: Implication for the formation of Tournaisian oil pools in the Volga–Ural anteklise, *Geol., Geofiz. Razrab. Neft. Gaz. Mestorozhd.*, 2006, no. 9, pp. 31–38.
- [11] Purser, B.H., Tucker, M.E., and Zenger, D.H., Problems, progress and future research concerning dolomites and dolomitization, in *Dolomites. A Volume in Honour of Dolomieu*. IAS Spec. Publ. N 21, Oxford: Blackwell, 1994.
- [12] Shvetsov, M.S., *Petrografiya osadochnykh porod (Petrography of Sedimentary Rocks)*, Moscow: Gosgeoltekhizdat, 1958.
- [13] Thompson, G., A geochemical study of some lithified carbonate sediments from the deep-sea, *Geochim. Cosmochim. Acta*, 1972, vol. 36, no. 11, pp. 1237–1253.
- [14] Tucker, V.E. and Wright, V.P., *Carbonate Sedimentology*, Oxford: Blackwell Science, 1990.
- [15] Warren, J, Dolomite: occurrence, evolution and economically important associations, *Earth Sci. Rev.*, 2000, vol. 52, pp. 1–81.